

State of the Art Updated Review of Tools for Assessment of Pain in Nonverbal Older Adults

Abbey, J., Piller, N., De Bellis, A., Esterman, A., Parker, D., Giles, L., et al. (2004). The abbey pain scale: A 1-minute numerical indicator for people with end-stage dementia. *International Journal of Palliative Nursing, 10*(1), 6-13.

AGS Panel on Persistent Pain in Older Persons. (2002). The management of persistent pain in older persons. *Journal of the American Geriatrics Society, 50*(6 Suppl), S205-24.

Baiardi, J., Parzuchowski, J., Kosik, C., Ames, T., Courtney, N., & Locklear, J. (2002). *Examination of the reliability of the FLACC Pain Assessment tool with cognitively impaired elderly*. Paper presented at the Annual National Conference of Gerontological Nurse Practitioners, Chicago, Ill.

Basler, H. D., Huger, D., Kunz, R., Luckmann, J., Lukas, A., Nikolaus, T., et al. (2006). Assessment of pain in advanced dementia. construct validity of the german PAINAD. [Beurteilung von Schmerz bei Demenz (BESD). Untersuchung zur Validitat eines Verfahrens zur Beobachtung des Schmerzverhaltens] *Schmerz (Berlin, Germany), 20*(6), 519-526.

Bauer, C., Lahjibi-Paulet, H., Somme, D., Onody, P., Saint Jean, O., & Gisselbrecht, M. (2007). Tolerability of an equimolar mix of nitrous oxide and oxygen during painful procedures in very elderly patients. *Drugs & Aging, 24*(6), 501-507.

Burns, N., & Grove, S. K. (2001). *The Practice of Nursing Research: Conduct, Critique, & Utilization* (4th ed.). Philadelphia: W.B. Saunders Company.

Cervo, F. A., Raggi, R. P., Bright-Long, L. E., Wright, W. K., Rows, G., Torres, A. E., et al. (2007). Use of the certified nursing assistant pain assessment tool (CPAT) in nursing

home residents with dementia. *American Journal of Alzheimer's Disease and Other Dementias*, 22(2), 112-119.

Cohen-Mansfield, J. (2006). Pain assessment in noncommunicative elderly persons--PAIN. *The Clinical Journal of Pain*, 22(6), 569-575.

Cohen-Mansfield, J., & Lipson, S. (2008). The utility of pain assessment for analgesic use in persons with dementia. *Pain*, 134(1-2), 16-23.

Cornali, C., Franzoni, S., Gatti, S., & Trabucchi, M. (2006). Diagnosis of chronic pain caused by osteoarthritis and prescription of analgesics in patients with cognitive impairment. *Journal of the American Medical Directors Association*, 7(1), 1-5.

Costardi, D., Rozzini, L., Costanzi, C., Ghianda, D., Franzoni, S., Padovani, A., et al. (2007). The Italian version of the pain assessment in advanced dementia (PAINAD) scale. *Archives of Gerontology and Geriatrics*, 44(2), 175-180.

Davies, E., Male, M., Reimer, V., & Turner, M. (2004). Pain assessment and cognitive impairment: Part 2. *Nursing Standard (Royal College of Nursing (Great Britain))*: 1987, 19(13), 33-40.

Davies, E., Male, M., Reimer, V., Turner, M., & Wylie, K. (2004). Pain assessment and cognitive impairment: Part 1. *Nursing Standard (Royal College of Nursing (Great Britain))*: 1987, 19(12), 39-42.

DeWaters, T., Faut-Callahan, M., McCann, J. J., Paice, J. A., Fogg, L., Hollinger-Smith, L., et al. (2008). Comparison of self-reported pain and the PAINAD scale in hospitalized cognitively impaired and intact older adults after hip fracture surgery. *Orthopaedic Nursing / National Association of Orthopaedic Nurses*, 27(1), 21-28.

Doloplus Group Website. <http://www.doloplus.com>

Feldt, K. S. (1996). Treatment of pain in cognitively impaired versus cognitively intact post hip fractured elders. (Doctoral dissertation, University of Minnesota, 1996). Dissertation Abstracts International, 57-09B, 5574.

Feldt, K. S. (2000). The checklist of nonverbal pain indicators (CNPI). *Pain Management Nursing : Official Journal of the American Society of Pain Management Nurses*, 1(1), 13-21.

Feldt, K. S. (2000). Improving assessment and treatment of pain in cognitively impaired nursing home residents. *Annals of Long Term Care*, 8(9), 36-42.

Fisher, S. E., Burgio, L. D., Thorn, B. E., Allen-Burge, R., Gerstle, J., Roth, D. L., et al. (2002). Pain assessment and management in cognitively impaired nursing home residents: Association of certified nursing assistant pain report, minimum data set pain report, and analgesic medication use. *Journal of the American Geriatrics Society*, 50(1), 152-156.

Fuchs-Lacelle, S., & Hadjistavropoulos, T. (2004). Development and preliminary validation of the pain assessment checklist for seniors with limited ability to communicate (PACSLAC). *Pain Management Nursing : Official Journal of the American Society of Pain Management Nurses*, 5(1), 37-49.

Fuchs-Lacelle, S., & Hadjistavropoulos, T. (2005). A checklist for pain assessment in LTC -- PACSLAC: Pain assessment checklist for seniors with limited ability to communicate. *Canadian Nursing Home*, 16(4), 4-7.

Fuchs-Lacelle, S., Hadjistavropoulos, T. & Lix, L. (in press). Pain assessment as intervention: A study of older adults with severe dementia. *Clinical Journal of Pain*, xx ,xxx. (in press)

Grbich, C., Maddocks, I., Parker, D., Brown, M., Willis, E., Piller, N., et al. (2005).

Identification of patients with noncancer diseases for palliative care services. *Palliative & Supportive Care*, 3(1), 5-14.

Herr, K., Bjoro, K., & Decker, S. (2006). Tools for assessment of pain in nonverbal older adults with dementia: A state-of-the-science review. *Journal of Pain and Symptom Management*, 31(2), 170-192.

Holen, J. C., Saltvedt, I., Fayers, P. M., Bjornnes, M., Stenseth, G., Hval, B., et al. (2005). The norwegian doloplus-2, a tool for behavioural pain assessment: Translation and pilot-validation in nursing home patients with cognitive impairment. *Palliative Medicine*, 19(5), 411-417.

Holen, J. C., Saltvedt, I., Fayers, P. M., Hjermstad, M. J., Loge, J. H., & Kaasa, S. (2007). Doloplus-2, a valid tool for behavioural pain assessment? *BMC Geriatrics*, 7, 29.

Horgas, A. L., Nichols, A. L., Schapson, C. A., & Vietes, K. (2007). Assessing pain in persons with dementia: Relationships among the non-communicative patient's pain assessment instrument, self-report, and behavioral observations. *Pain Management Nursing : Official Journal of the American Society of Pain Management Nurses*, 8(2), 77-85.

Hurley, A. C., Volicer, B. J., Hanrahan, P. A., Houde, S., & Volicer, L. (1992). Assessment of discomfort in advanced alzheimer patients. *Research in Nursing & Health*, 15(5), 369-377.

Husebo, B. S., Strand, L. I., Moe-Nilssen, R., Borgehusebo, S., Aarsland, D., & Ljunggren, A. E. (2008). Who suffers most? dementia and pain in nursing home patients: A cross-sectional study. *Journal of the American Medical Directors Association*, 9(6), 427-433.

Husebo, B. S., Strand, L. I., Moe-Nilssen, R., Husebo, S. B., Snow, A. L., & Ljunggren, A. E.

(2007). Mobilization-observation-behavior-intensity-dementia pain scale (MOBID): Development and validation of a nurse-administered pain assessment tool for use in dementia. *Journal of Pain and Symptom Management, 34*(1), 67-80.

Hutchison, R. W., Tucker, W. F., Jr, Kim, S., & Gilder, R. (2006). Evaluation of a behavioral assessment tool for the individual unable to self-report pain. *The American Journal of Hospice & Palliative Care, 23*(4), 328-331.

Jacobson, S.F.(1997). Evaluating instruments for use in clinical nursing research. In M. Frank-Stromborg & S.J. Olsen (eds.). *Instruments for Clinical Health Care Research*. Boston, MA: Jones and Bartlett Publishers.

Jones, K. R., Fink, R., Hutt, E., Vojir, C., Pepper, G. A., Scott-Cawiezell, J., et al. (2005). Measuring pain intensity in nursing home residents. *Journal of Pain and Symptom Management, 30*(6), 519-527.

Kovach, C. R., Weissman, D. E., Griffie, J., Matson, S., & Muchka, S. (1999). Assessment and treatment of discomfort for people with late-stage dementia. *Journal of Pain and Symptom Management, 18*(6), 412-419.

Kovach, C. R., Noonan, P. E., Griffie, J., Muchka, S., & Weissman, D. E. (2001). Use of the assessment of discomfort in dementia protocol. *Applied Nursing Research : ANR, 14*(4), 193-200.

Kovach, C. R., Noonan, P. E., Griffie, J., Muchka, S., & Weissman, D. E. (2002). The assessment of discomfort in dementia protocol. *Pain Management Nursing : Official Journal of the American Society of Pain Management Nurses, 3*(1), 16-27.

- Lane, P., Kuntupis, M., MacDonald, S., McCarthy, P., Panke, J. A., Warden, V., et al. (2003). A pain assessment tool for people with advanced alzheimer's and other progressive dementias. *Home Healthcare Nurse, 21*(1), 32-37.
- Lefebvre-Chapiro, S. (2001). The Doloplus 2 scale - evaluating pain in the elderly. *European Journal of Palliative Care, 8*(5), 191-194.
- Leong, I. Y., Chong, M. S., & Gibson, S. J. (2006). The use of a self-reported pain measure, a nurse-reported pain measure and the PAINAD in nursing home residents with moderate and severe dementia: A validation study. *Age and Ageing, 35*(3), 252-256.
- Merkel, S. I., Voepel-Lewis, T., Shayevitz, J. R., & Malviya, S. (1997). The FLACC: A behavioral scale for scoring postoperative pain in young children. *Pediatric Nursing, 23*(3), 293-297.
- Miller, J., Neelon, V., Dalton, J., Ng'andu, N., Bailey, D., Jr., Layman, E., et al. (1996). The assessment of discomfort in elderly confused patients: a preliminary study. *Journal of Neuroscience Nursing, 28*(3), 175-182.
- Morello, R., Jean, A., Alix, M., Sellin-Peres, D., & Fermanian, J. (2007). A scale to measure pain in non-verbally communicating older patients: The EPCA-2 study of its psychometric properties. *Pain, 133*(1-3), 87-98.
- Norbeck, J. S. (1985). What constitutes a publishable report of instrument development? *Nursing Research, 34*(6), 380-382.
- Nygaard, H. A., & Jarland, M. (2006). The checklist of nonverbal pain indicators (CNPI): Testing of reliability and validity in norwegian nursing homes. *Age and Ageing, 35*(1), 79-81.

- Pautex, S., Herrmann, F. R., Michon, A., Giannakopoulos, P., & Gold, G. (2007). Psychometric properties of the doloplus-2 observational pain assessment scale and comparison to self-assessment in hospitalized elderly. *The Clinical Journal of Pain, 23*(9), 774-779.
- Pautex, S., Michon, A., Guedira, M., Emond, H., Le Lous, P., Samaras, D., et al. (2006). Pain in severe dementia: Self-assessment or observational scales? [references]. *Journal of the American Geriatrics Society, 54*(7), 1040-1045.
- Pickering, G., Jourdan, D., & Dubray, C. (2006). Acute versus chronic pain treatment in alzheimer's disease. *European Journal of Pain (London, England), 10*(4), 379-384.
- Porzsolt, F., Kojer, M., Schmidl, M., Greimel, E. R., Sigle, J., Richter, J., et al. (2004). A new instrument to describe indicators of well-being in old-old patients with severe dementia--the vienna list. *Health and Quality of Life Outcomes, 2*, 10.
- Regnard C., Matthews D., Gibson L., Clarke C. & Watson B. (2003). Difficulties in identifying distress and its causes in people with severe communication problems. *International Journal of Palliative Nursing 9*, 173-6.
- Regnard, C., Reynolds, J., Watson, B., Matthews, D., Gibson, L., & Clarke, C. (2007). Understanding distress in people with severe communication difficulties: Developing and assessing the disability distress assessment tool (DisDAT). *Journal of Intellectual Disability Research, 51*, 277-292.
- Schuler, M. S., Becker, S., Kaspar, R., Nikolaus, T., Kruse, A., & Basler, H. D. (2007). Psychometric properties of the german "pain assessment in advanced dementia scale" (PAINAD-G) in nursing home residents. *Journal of the American Medical Directors Association, 8*(6), 388-395.

Snow, A.L., Hovanec, L., Passano, J., Brandt, J. (2001). Development of a pain assessment instrument for use with severely demented patients. Poster session presented at the Annual Meeting of the American Psychological Association. Washington, DC.

Snow, A. L., Weber, J. B., O'Malley, K. J., Cody, M., Beck, C., Bruera, E., et al. (2004). NOPPAIN: A nursing assistant-administered pain assessment instrument for use in dementia. *Dementia and Geriatric Cognitive Disorders*, 17(3), 240-246.

Snow, A.L., Cody, M., Beck, C., Bruera, E., O'Malley, K.J. Aston, C., El-Issa, L., Kunik, M.E. (draft article, 2004). Nursing assistants as pain screeners for nursing home residents with severe dementia: In vivo validation of the NOPPAIN.

Stevenson, K. M., Brown, R. L., Dahl, J. L., Ward, S. E., & Brown, M. S. (2006). The discomfort behavior scale: A measure of discomfort in the cognitively impaired based on the minimum data set 2.0. *Research in Nursing & Health*, 29(6), 576-587.

Stolee, P., Hillier, L. M., Esbaugh, J., Bol, N., McKellar, L., & Gauthier, N. (2005). Instruments for the assessment of pain in older persons with cognitive impairment. *Journal of the American Geriatrics Society*, 53(2), 319-326.

Tsai, P. F., Beck, C., Richards, K. C., Phillips, L., Roberson, P. K., & Evans, J. (2008). The pain behaviors for osteoarthritis instrument for cognitively impaired elders (PBOICIE). *Research in Gerontological Nursing*, 1(2), 116-122.

van Herk, R., van Dijk, M., Baar, F. P., Tibboel, D., & de Wit, R. (2007). Observation scales for pain assessment in older adults with cognitive impairments or communication difficulties. *Nursing Research*, 56(1), 34-43.

- van Iersel, T., Timmerman, D., & Mullie, A. (2006). Introduction of a pain scale for palliative care patients with cognitive impairment. *International Journal of Palliative Nursing*, 12(2), 54-59.
- Villanueva, M. R., Smith, T. L., Erickson, J. S., Lee, A. C., & Singer, C. M. (2003). Pain assessment for the dementing elderly (PADE): Reliability and validity of a new measure. *Journal of the American Medical Directors Association*, 4(1), 1-8.
- Warden, V., Hurley, A. C., & Volicer, L. (2003). Development and psychometric evaluation of the pain assessment in advanced dementia (PAINAD) scale. *Journal of the American Medical Directors Association*, 4(1), 9-15.
- Wary, B., & Doloplus, C. (1999). Doloplus-2, a scale for pain measurement. [Doloplus-2, une échelle pour évaluer la douleur] *Soins.Gerontologie*, (19)(19), 25-27.
- Zwakhalen, S. M., Hamers, J. P., Abu-Saad, H. H., & Berger, M. P. (2006). Pain in elderly people with severe dementia: A systematic review of behavioural pain assessment tools. *BMC Geriatrics*, 6, 3.
- Zwakhalen, S. M., Hamers, J. P., & Berger, M. P. (2007). Improving the clinical usefulness of a behavioural pain scale for older people with dementia. *Journal of Advanced Nursing*, 58(5), 493-502.
- Zwakhalen, S. M. G., Hamers, J. P. H., & Berger, M. P. F. (2006). The psychometric quality and clinical usefulness of three pain assessment tools for elderly people with dementia. *Pain*, 126(1-3), 210-220.
- Young, D. M. (2001). Pain in institutionalized elders with chronic dementia. *The University of Iowa ** Ph.D*, 169.